

Historical zooplankton composition indicates eutrophication stages in a Neotropical aquatic system: the case of Lake Amatitlán, Central America

Sarahi Jaime ^{1*}, Adrián Cervantes-Martínez ¹, Martha A. Gutiérrez-Aguirre ¹, Eduardo Suárez-Morales ², Julio R. Juárez-Pernillo ³, Elena M. Reyes-Solares ³, and Victor H. Delgado-Blas ¹

- ¹ Universidad de Quintana Roo (UQROO), Avenida Andrés Quintana Roo. Col. San Gervasio, Cozumel 77600, Quintana Roo, Mexico; 1518305@uqroo.mx; adcervantes@uqroo.edu.mx, margutierrez@uqroo.edu.mx
² El Colegio de la Frontera Sur (ECOSUR), Avenida Centenario Km 5.5, Chetumal 77014, Quintana Roo, Mexico; esuarez@ecosur.mx.
³ Autoridad para el Manejo Sustentable de la cuenca del lago de Amatitlán (AMSA); Kilómetro 22 CA-9, Bárcenas Villanueva, 6624-1700, Guatemala; jjpernillo@gmail.com, elena1reyes@gmail.com.
* Correspondence: 1518305@uqroo.mx

Abstract: This paper presents a study of freshwater zooplankton biodiversity, deemed as a reliable indicator of water quality. The Guatemalan Lake Amatitlán, currently used as a water source, has shown signs of progressive eutrophication, with perceptible variations of the local zooplankton diversity. Biotic and abiotic parameters were determined at four sites of Lake Amatitlán (i. e., Este Centro, Oeste Centro, Bahía Playa de Oro and Michatoya) in 2016 and 2017. The local composition, species richness and abundance of zooplankton, and the system environmental parameters were analyzed during both years surveyed. Biological data suggesting eutrophication of this tropical system were obtained, including a high rotifer abundance (11 species: the rotifers *Brachionus havanaensis* (109 ind L⁻¹) and *Keratella americana* (304 ind L⁻¹) were the most abundant species in this lake). The presumably endemic diaptomid copepod species *Mastigodiaptomus amatitlanensis*, was absent in our samples, but we report the unprecedented occurrence of two Asian cyclopoid copepods (i. e., *Thermocyclops crassus*, *Mesocyclops thermocyclopoides*) for Lake Amatitlán and Guatemala. The presence of larger zooplankters like adults and immature copepods (*Arctodiaptomus dorsalis*) and cladocerans (*Ceriodaphnia* sp.) at site “Este Centro” indicates a relatively healthy zooplankton community and represents a focal point for managing the conservation of this lake.

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *Diversity* **2021**, *13*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Lastname

Received: date
Accepted: date
Published: date

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: conservation; eutrophication; exotic species; tropical lakes; zooplankton

1. Introduction

The knowledge of zooplankton in the Neotropical region is growing with fragmented studies. Therefore, it is likely that the species richness of zooplanktonic taxa is underestimated because of the presumably high diversity and the scarcity of zooplankton taxonomists [1-3]. In addition, the progressive destruction of aquatic habitat and the progressive spread of exotic species threaten native biodiversity, ecosystem health, and environmental services.

The zooplankton community and abundance are closely linked to the trophic state of the water system, for this reason, its diversity has been deemed as an indicator of water quality [4]. In eutrophicated systems (at tropical and temperate latitudes), the dominance of microzooplankton is common, compared to larger organisms, due to the increased availability of food and water conditions [5,6]

For 4 decades, the Guatemalan Lake Amatitlán has shown signs of progressive eutrophication related to anthropic factors (i. e. peripheral population growth and

urbanization, intensive use of water for agricultural irrigation), thus, promoting the advancement towards eutrophication related to the input of nearly 50% of the untreated residual urban and industrial waters from Guatemala City [7–10]. Because of this, some actions have been proposed to address this problem, either from the governmental level (i.e., Autoridad para el Manejo Sustentable de la cuenca del lago Amatitlán, AMSA 1996) and from descriptive studies of the lake involving the lake zooplankton biodiversity, like those by Basterrechea-Díaz (1997) [7] and Brandorff (2012) [11]; however, studies related with tropical epicontinental waterbodies have been more focused on environmental factors rather than biological or community attributes or general limnology [12,13]; thus, the zooplankton biodiversity in Guatemala remains largely unknown [14], with only a few studies in Guatemalan lakes [15, 16]. Most studies in Lake Amatitlán and Guatemala, are more focused on current data instead of historical analysis.

Based on the analysis of both, historical and current data of zooplankton biodiversity and environmental conditions of Lake Amatitlán, we present information on the zooplankton distribution, species richness, abundance, and its relation with successive changes of its trophic state.

2. Materials and Methods

2.1. Study sites and sampling methods

Lake Amatitlán is the fourth largest lake in Guatemala, Central America, and one of the most emblematic waterbodies of this country. This lake is a warm monomictic waterbody in the highland of Guatemala, located at an altitude of 1,186 meters above sea level (m.a.s.l.), with an area of 15.2 km² and 11 km length and a maximum depth of 23 m. Its formation is originated from volcanic activity of Pacaya, Fuego, and Agua in the late quaternary. [10,14,17].

Four sampled sites were considered: Este Centro (EC), Oeste Centro (OC), Bahía Playa de Oro (BPO) and Michatoya (MICH) to analyze the zooplankton species that inhabit the east and west regions of Lake Amatitlán (Figure 1). The latter two sites (BPO and MICH) are in the runoff of Villalobos and Michatoya rivers respectively [14]. Water samples for biotic and abiotic variables were collected for 2016 and 2017 in the rainy (May–October) and dry seasons (November–April).




Figure 1. Location of Lake Amatitlán and sampling points with information of biotic collection methods. Water filtered, vertical and horizontal trawls as defined by Cervantes-Martínez & Gutiérrez-Aguirre (2015) [2].

2.1.1. Species richness

Zooplankton samples ($n = 8$) were collected by vertical and horizontal trawls with a 45 μm plankton net between 1-22 meters depth, to ensure representative samples to evaluate the species richness in the lake, since it is well known that zooplankton tends to have vertical and horizontal migrations [2].

2.1.2. Species abundance and abiotic variables

To estimate the zooplankton abundance, a known volume of water between 30 to 100 liters was filtered through a 45 μm zooplankton net. The water was determined with a 2.1 L⁻¹ capacity Van Dorn bottle [2, 18]. Species abundance was determined by the account of two main groups: Rotifera and Copepoda, present in 3 aliquots of 1 ml each from the filtered samples, then the data were standardized as individuals per liter (ind L⁻¹) in each sampled site [19].

Abiotic variables were measured *in situ* monthly for both years of study and in all the water column, with the multiparametric probes WTW Cond 197i, WTW Oxi 1970i, and HACH HQ for water temperature ($^{\circ}\text{C}$), pH, oxygen concentration O₂ (mg L⁻¹), total dissolved solids (mg L⁻¹), and conductivity ($\mu\text{S cm}^{-1}$). With the actual environmental, richness and zooplankton abundance data, a description about the trophic state of Lake Amatitlán, was proposed.

2.2. Historical and actual records of zooplankton and environmental parameters analysis

Specific classification of Rotifera, Cladocera and Copepoda of recently collected samples (collected in 2016, and 2017) was done according to Koste (1978) [20], Fontaneto & De Smet (2015) [21], Elías-Gutiérrez *et al.* (2008) [22] and Suárez-Morales *et al.* (2020) [23].

The presence/absence of the current zooplankton inventory was compared with previous surveys by Juday (1915) [24], Basterrechea-Díaz (1997) [7], and the record of copepods from the previous surveys of Wilson (1941) [25], and Brandorff (2012) [11], in order to analyze the historical composition of zooplankton of Amatitlán lake.

Historical environmental data recorded by Juday (1915) [24], Brezonik & Fox (1974) [26], Basterrechea-Díaz (1997) [7], and Ellenberg (2014) [27] were compared with the current data surveyed in this study.

3. Results

3.1.1. Species richness

A total of 15 species of zooplankters including rotifers and crustaceans were found in the lake for 2016-2017 (**Table 1**); rotifers showed the highest species richness (80% of zooplankton species recorded) while copepods represented 20% of all zooplankton species in the lake.

We provide the first record of two cyclopoid exotic species (*Mesocyclops thermocyclopoides* and *Thermocyclops crassus*) for Lake Amatitlán and Guatemala. The endemic calanoid copepod *Mastigodiatomus amatitlanensis*, was absent in our current survey and the record of *Arctodiatomus dorsalis* in Amatitlán was confirmed here. Cladoceran crustaceans were very scarce in our samples, a single specimen of *Ceriodaphnia* sp. was observed. The Brachionidae was the family with the highest species richness among rotifers in 2016 and 2017 (**Table 1**).

Nowadays, the east region (site EC) of Lake Amatitlán had the highest species richness in the lake (14 species), compared with western region (9 species including the exotic *T. crassus*). The largest zooplankters of the lake, including the cladoceran *Ceriodaphnia*, (~2

mm) [21], the calanoid copepod *A. dorsalis* and the cyclopoid copepod *M. thermocyclopoides* occurred in eastern region.

Our revision of the zooplankton community (Table 1) indicates that the historical data presented a great microcrustacean richness with the record of 8 cladoceran species (*Daphnia* sp., *D. hyalina*, *Ceriodaphnia* sp. *C. lacustris*, *C. pulchella*, *Bosmina* sp., *B. longirostris*, and *Chydorus sphaericus*) and the 3 calanoid copepods: *A. dorsalis*, *Mastigodiptomus albuquerqueensis*, and the endemic *M. amatitlanensis*. The historical record of rotifers had the lowest species richness including three monogonont species. In our survey, the rotifer species richness increased significantly with 12 species not hitherto reported from the lake, including the record of organisms from the Subclass Bdelloidea.

Table 1. Current and historical records of zooplankton species richness in Lake Amatitlán. The currently recorded species in columns 1) EC, 2) OC, 3) BPO and 4) MICH. Historical records in columns 5-8, following data by Brandorff (2012) [11] Basterrechea-Díaz (1997) [7]; Wilson (1941) [25]; and Juday (1915) [24], respectively. Presence (x), absence (-), new records (*).

Species	Current Data				Historical Data			
	1	2	3	4	5	6	7	8
Phylum: Rotifera								
Monogononta: Ploimida								
Family: Epiphanidae Haring, 1913								
<i>Epiphanes macroura</i> Barrois & Daday, 1894*	x	x	-	-	-	-	-	-
Family: Brachionidae Ehrenberg, 1838								
<i>Anuraeopsis fissa</i> (Gosse, 1851)*	x	-	-	-	-	-	-	-
<i>Brachionus angularis</i> (Gosse, 1851) *	x	x	x	x	-	-	-	-
<i>B. calyciflorus</i> Pallas, 1766 *	x	x	x	x	-	-	-	-
<i>B. plicatilis</i> Müller, 1786 *	x	-	-	x	-	-	-	-
<i>B. havanaensis</i> Rousselet, 1911*	x	x	x	x	-	-	-	-
<i>Keratella</i> sp.	-	-	-	-	-	x	-	-
<i>K. americana</i> Carlin, 1943*	x	x	x	x	-	-	-	-
<i>K. cochlearis</i> (Gosse, 1851)	-	-	-	-	-	-	-	x
Family: Trichocercidae Haring, 1913								
<i>Trichocerca</i> cf. <i>longiseta</i> (Schrank, 1802)*	-	x	x	x	-	-	-	-
<i>T. pusilla</i> (Lauterborn, 1898)*	-	x	x	x	-	-	-	-
Family: Asplanchnidae Eckstein, 1883								
<i>Asplanchna sieboldi</i> (Leydig, 1854)*	x	x	x	-	-	-	-	-
Flosculariaceae: Family: Trochosphaeridae Haring, 1913								
<i>Filinia longiseta</i> (Ehrenberg, 1834)	x	x	x	x	-	-	-	x
<i>F. terminalis</i> (Plate, 1886) *	x	x	x	-	-	-	-	-
Subclass: Bdelloidea*	x	-	-	-	-	-	-	-
Superclass: Crustacea								
Brachiopoda: Cladocera: Anomopoda								
Family: Daphniidae Straus, 1820								
<i>Daphnia</i> sp.	-	-	-	-	-	x	-	-
<i>D. hyalina</i> Leydig, 1860	-	-	-	-	-	-	-	x
<i>Ceriodaphnia</i> sp.	x	-	-	-	-	x	-	x
<i>C. lacustris</i> Birge, 1893	-	-	-	-	-	-	-	x
<i>C. pulchella</i> Sars, 1862	-	-	-	-	-	-	-	x
Family: Bosminidae Sars, 1865								
<i>Bosmina</i> sp.	-	-	-	-	-	x	-	-
<i>Bosmina longirostris</i> O. F. Müller, 1776	-	-	-	-	-	-	-	x
Family: Chydoridae Stebbing, 1902								
<i>Chydorus sphaericus</i> (O.F. Müller, 1785)	-	-	-	-	-	-	-	x
Copepoda: Calanoida								
Family: Diaptomidae G.O. Sars, 1932								
Subfamily: Diaptominae Kiefer, 1932								

<i>Arctodiaptomus dorsalis</i> (Marsh, 1907)	x	-	-	-	x	-	-	-
<i>Mastigodiaptomus albuquerquensis</i> (Herrick, 1895)	-	-	-	-	-	-	-	x
<i>M. amatitlanensis</i> (Wilson, 1941)	-	-	-	-	-	-	x	-
Copepoda: Cyclopoida								
Family: Cyclopidae Kiefer, 1927								
Subfamily: Cyclopinae Kiefer, 1927								
<i>Thermocyclops crassus</i> (Fischer, 1853)*	-	x	-	-	-	-	-	-
<i>Mesocyclops thermocyclopoides</i> Harada, 1931*	x	-	-	-	-	-	-	-
Subfamily: Eucyclopinae Kiefer, 1927								
<i>Eucyclops serrulatus</i> (Fischer, 1851)	-	-	-	-	-	-	-	x
Nauplii	x	x	x	x	-	x	-	x
Juvenile Cyclopoid	x	x	x	x	-	-	-	-
Juvenile Calanoid	x	x	x	x	-	-	-	-

3.1.2. Species Abundance

In this study, the total rotifer abundance was 522.7 ind L⁻¹. Rotifers represent the most abundant group in the lake; their numerical abundance is considerably higher than that recorded for copepods, including immature stages (7.1 ind L⁻¹). Cladocerans were almost absent from our samples.

Species with the highest abundance at all sites were the rotifers *B. havanaensis* (109 ind L⁻¹) and *K. americana* (304 ind L⁻¹), with a considerably lower abundance in the eastern area (9.3 and 121.8 ind L⁻¹ respectively). Species of the family Brachionidae were the most abundant mainly in western region (sites OC, BPO and MICH), whereas the lowest abundance of rotifers occurred in the eastern region (site EC) (see **Table 2**).

Table 2. Abundance (ind L⁻¹) calculated from zooplankton samples for all the studied points of Lake Amatitlán in 2017.

Species	Abundance (ind L ⁻¹)			
	EC	OC	BPO	MICH
<i>Brachionus angularis</i>	0.00	0.00	71.56	1.87
<i>Brachionus calyciflorus</i>	0.70	0.70	85.56	4.67
<i>Brachionus plicatilis</i>	0.00	0.47	0.00	0.00
<i>Trichocerca cf. longiseta</i>	0.00	0.00	14.00	12.60
<i>Trichocerca pusilla</i>	0.00	8.40	13.22	11.20
<i>Asplanchna sieboldi</i>	1.40	1.17	2.33	0.93
<i>Filinia longiseta</i>	0.00	1.40	28.00	13.07
<i>Filinia terminalis</i>	8.17	49.23	35.00	60.20
<i>Brachionus havanaensis</i>	9.33	153.53	108.89	165.20
<i>Keratella americana</i>	121.80	432.60	265.22	408.33
Nauplii	3.50	2.57	3.11	1.40
Juvenile Cyclopoid	5.83	1.63	1.56	0.93
Juvenile Calanoid	2.80	0.47	3.89	0.47
<i>M. thermocyclopoides</i>	0.23	0.00	0.00	0.00

The local copepod abundance was represented mainly by nauplii and juvenile stages of Calanoida and Cyclopoida (average= 2.6, 2.5 and 1.9 ind L⁻¹, respectively), values resembling those recorded for the Rotifera like *B. plicatilis* (1.1 ind L⁻¹) and *A. sieboldi* (2.3 ind L⁻¹) in all the study sites, compared to adult copepods, where the abundance of the adult *M. thermocyclopoides* present only in EC, was 0.23 ind L⁻¹.

3.2. Environmental variables

Environmental variables values in both analyzed years, in general, presented basic pH values, (> 8 ± 0.33), dissolved oxygen showed an average of 4.76 ± 5.21 and 4.65 ± 4.92 mg L⁻¹, whereas temperature averaged 24 ± 1.31 °C, conductivity presented average values of 655.95 ± 59.52 and 678.23 ± 68.29 µS cm⁻¹, finally TDS showed average values of 339.99 ± 47.35 and 341.43 ± 30.30 mg L⁻¹ respectively (Table 3).

The historical data presented in **Table 3** show pH with slightly neutral values in 1969 to 1985-1995, whereas in the first two decades of the XXI century, the pH increased to reach clearly basic values, over 8. The water temperature changed along the time, 19.86 °C in 1910 to 24.23 °C in 2017. Conductivity and Total Dissolved Solids decreased in average 18.29 and 44.10%, respectively.

Table 3. Historical and current environmental mean data of the water column recorded by previous surveys and this study. [24] Juday (1915), [26] Brezonik & Fox (1974), [7] Basterrechea-Díaz (1997), [27] Ellenberg (2014). ND: No Data available.

Environmental variables	1910 [24]	1969 [26]	1985-1995 [7]	2008-2013 [27]	2016 ***	2017 ***
pH	ND	7.70	7.75	8.69	8.26	8.33
Water temperature (°C)	19.86	ND	22.75	25.00	24.46	24.23
Conductivity ($\mu\text{S cm}^{-1}$)	ND	830	802	682	655.95	678.23
TDS (mg L^{-1})	ND	ND	610	ND	339.99	341.43
Dissolved Oxygen O_2 (mg L^{-1})	4.74*	8.40**	4.20	8.90	4.76	4.65

* Data originally recorded in Cubic Centimeters per liter of water.

** Original data recorded at surface.

*** Data recorded in this study.

4. Discussion

The environmental parameters surveyed in this study can show the progressive eutrophication on Lake Amatitlán, according to the historical data recorded by authors like Juday (1915) [24], Brezonik & Fox (1974) [26] Basterrechea-Díaz (1997) [7], and Ellenberg (2014) [27]. The historical change in environmental and biological variables could reveal strong evidence of the current eutrophication of this lake. For instance, the observed changes of pH values: an average of 8.26 and 8.33 in 2016-17 differ in contrast from the values recorded in 1969 (7.70) [26], 1985-1995 (7.75) [7], and 2008 (9.3) [17].

The basic pH and the high concentration of dissolved oxygen at the surface promoted an increase of microzooplankters, like rotifers (especially *B. havanaensis* and *K. americana*) and a decrease of larger species like cladocerans and adult copepods, indicators of the system trophic state *per se*. Similar conditions have been recorded in American eutrophicated subtropical and tropical water bodies [4,28,29] and also in other water bodies (i. e., temperate coastal water bodies) in which the replacement of larger copepod with smaller ones has been reported resulting from the eutrophication process [6].

Recently, the phytoplankton blooming has been described as a consequence of this eutrophication progress in Lake Amatitlán, presenting a high concentration mainly in *Microcystis* sp. and *Dolichospermum* sp. cyanobacteria preceded by the diatom algae *Nitzschia* sp. at the surface of the lake [9], which in turn allows herbivorous zooplankters like brachionid rotifers to become dominant organisms in eutrophicated epicontinental waterbodies [20]. In earlier studies on Lake Amatitlán, the zooplankton community was largely dominated by cladocerans and copepods. In 1915 [24], zooplankton had a widely different composition compared to our results: rotifers were then the less abundant zooplankton group in the lake (0.3 ind L^{-1}), preceded by copepods (11.6 ind L^{-1}) and cladocerans, the most abundant zooplankton group at that time (14.4 ind L^{-1}). The system trophic state is also related to the zooplankters body size: a stronger level of eutrophication is frequently expressed by a greater abundance and species richness of microzooplankters like small rotifers [4,6,28,29]. A possible explanation of the local absence or scariness of larger zooplankters (i. e. *Ceriodaphnia* sp., adult cyclopoid and calanoid copepods, including *M. amatitlanensis*), could result from the competition for available food [5], eventually explaining the strong dominance of small brachionid herbivorous rotifers like *B. havanaensis* and *K. americana*.

The presence and high abundance of these latter species, together with another species of *Brachionus* and *Keratella* at the east region of Amatitlán suggest that eutrophic conditions make food available for these microphagous species [30].

In the case of *A. dorsalis*, this species is widespread in America [31] and has been recorded as an invasive exotic copepod in Asiatic waterbodies [32,33]. The environmental conditions of Lake Amatitlán seem to be adequate for the development of this species because it shows a selective feeding on phytoplankton; thus, it frequently inhabits moderately to strongly eutrophic environments [31,32], like Amatitlán.

It is well known that many diaptomid copepods tend to have restricted distributional patterns and endemic distributions in neotropical lakes [34]. Then, the local absence of the endemic copepod *M. amatitlanensis* in this study could be another indication of the progressive eutrophication of Lake Amatitlán; because since its description by Wilson (1941) [25], this species has not been recorded in other regional studies (*i. e.* Elías-Gutiérrez et al., 2008 [35]; Brandorff, 2012 [11] and Gutiérrez-Aguirre, et al., 2020 [36]) It is probable that *M. amatitlanensis* occurs in other lakes of Guatemala (or Central America) and it is expected to be collected from adjacent systems. It is also probable that this species dwells at higher depths not easily reached by standard nets.

Our results showed a clear zonification; the eastern region (site EC) diverges from the other sites because of the absence of adjacent rivers (see **Figure 1**), its distance from the other sampling points (the closest site is OC, 7.04 km away) and its separation from other sites due to a train riel which divides the lake in two [14]. Therefore, the EC area has the best conservation status of the lake, precisely where we found the greatest species richness and the larger zooplankters, like the copepods *T. crassus* (average body length of 0.56–0.93 mm) [37], *M. thermocycloides* (0.78–0.89 mm) [38], and *A. dorsalis* (0.77–1.13 mm) [31] among them. Thus, it is convenient to consider EC as a potential conservation site since it has better environmental conditions for the conservation and preservation of zooplankton biodiversity.

On the other hand, we report the presence of two exotic cyclopoid copepod species for the Central American Lake Amatitlán and Guatemala country, *M. thermocycloides* and *T. crassus*. *M. thermocycloides* is a native species from Taiwan and well spread in Asia and Africa, and commonly widespread at tropical latitudes. This species have been recorded in lakes from South Mexico in epicontinental waterbodies from Chiapas state, Mexico, considered that their introduction may be related to anthropic factors (*i. e.* agriculture, and aquaculture) [37,38]. This is the second record of the invasion of these species in Central American countries, since they has been recorded before in Costa Rican water bodies by Collado et al. (1894) [39], and it is well known the ecological potential of *Mesocyclops* use as biocontrol of vector mosquitoes like *Aedes aegypti*, [40–42]. Therefore, its finding in Guatemalan lakes represents a source for mass culture of this copepod to be used as biocontrol.

Thermocyclops crassus is commonly spread at tropical latitudes in Africa, Australia and Asia, it was recorded in Laurentian great lakes in the United States of America [43], recorded for the first time in tropical lakes from Tabasco state, Mexico [37] and in small ponds of San José Province in Costa Rica [39]. Being a thermophilic species, *T. crassus* has a narrow temperature tolerance, being a thermophilic species [44], so it may be a local indicator of the temperature changes in the lake along time.

Finally, the physical, chemical, and biological conditions of the lake have clearly changed over time, from being a lake with oligotrophic characteristics to hypertrophic conditions in a relatively short period of time, (100 years, approximately) and allowed us to follow and describe the stages and speed of the eutrophication process of a large neotropical lake.

5. Conclusions

The historical analysis of zooplankton composition in the lake presented in this study reinforce the knowledge of its eutrophic state, suggesting a useful role of the zooplankton as bioindicators and made possible the visualization of the changes in its composition over time, showing the progressive trophic state towards to eutrophic or hypereutrophic conditions.

Probably the absence of the endemic species *M. amatitlanensis* is a warning sign regarding the accelerated loss of biodiversity and reinforces the idea that zooplankton is a great tool as a bioindicator of the health status for continental aquatic ecosystems, in both, tropical and temperate latitudes.

Further studies analyzing bottom sediments to search resting eggs of zooplankton in Lake Amatitlán and around it, can answer the question of the absence of *M. amatitlanensis*, where this type of knowledge is also scarce in inland aquatic systems of the region.

Finally, is convenient to consider the isolated site EC as a focal point for conservation since it presents better environmental conditions for the conservation and preservation of zooplankton biodiversity, due to the record of largest zooplankters found in this site.

Author Contributions: Conceptualization, S.J. and A.C.M.; methodology, S.J, A.C.M. M.A.G.A, E.S.M, J.R.J.P, E.M.R.S, and V.H.D.B; software, S.J; validation, S.J, A.C.M. M.A.G.A, E.S.M, J.R.J.P, E.M.R.S, and V.H.D.B.; formal analysis, S.J, A.C.M. M.A.G.A, E.S.M, J.R.J.P, E.M.R.S, and V.H.D.B; investigation, S.J, A.C.M. M.A.G.A, E.S.M, J.R.J.P, E.M.R.S, and V.H.D.B.; resources, J.R.J.P. and E.M.R.S; data curation, S.J.; writing—original draft preparation, S.J; writing—review and editing, S.J, A.C.M. M.A.G.A, E.S.M, J.R.J.P, E.M.R.S, and V.H.D.B, ; visualization, S.J. and A.C.M.; supervision, S.J and A.C.M; project administration, S.J, A.C.M. M.A.G.A, E.S.M, J.R.J.P, E.M.R.S, and V.H.D.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Council of Science and Technology (CONACYT) with the Investigator SNI-III Assistant Program, Grant Number 19812.

Institutional Review Board Statement: Zooplankton collected in this study were provided by AMSA in Guatemala, zooplankton is not under any protection by Guatemalan laws; therefore, no specific permits are required for this kind of field.

Acknowledgments: We gratefully acknowledge the invitation of Dr. Manuel Elías-Gutiérrez to contribute to this special issue of the Selected Papers from 1st International Electronic Conference on Biological Diversity, Ecology and evolution, and his valuable suggestions that made possible this research. We acknowledge AMSA for all the biological material and data bases that made possible the present study. Anonymous reviewers made valuable comments and suggestions. University of Quintana Roo, Cozumel with the investigation group vulnerability of continental and coastal water systems and the National Council of Science and Technology (CONACYT) with the Investigator SNI-III Assistant Program (Number 19812) gave financial support.

Conflicts of Interest: The authors declare no conflict of interest.

References

- José de Paggi, S. B.; Wallace, R.; Fontaneto, D.; Marinone, M. C. Phylum Rotifera. In *Thorpe and Covich's Freshwater Invertebrates*; Damborenea, C., Rogers, C. D., James, T., Eds.; Elsevier Science Publishing Co Inc, San Diego, United States, 2020; pp 145–200.
- Cervantes-Martínez, A.; Gutiérrez-Aguirre, M. A. Physicochemistry and Zooplankton of Two Karstic Sinkholes in the Yucatan Peninsula, Mexico. *J. Limnol.* **2015**, *74* (2), 382–393. <https://doi.org/10.4081/jlimnol.2014.976>.
- Suárez-Morales, E.; Reid, J. W.; Ilige, T.; Fiers, F. *Catálogo de Los Copépodos (Crustacea) Continentales de La Península de Yucatán, México*; Chetumal, Quintana Roo, México. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, Ciudad de México, México, 1996; 296 p.
- Gómez-Márquez, J. L.; Peña-Mendoza, B.; Guzmán-Santiago, J. L.; Gallardo-Pineda, V. Composición, Abundancia Del Zooplankton y Calidad de Agua En Un Microreservorio En El Estado de Morelos. *Hidrobiologica* **2013**, *23* (2), 227–240.
- Gama Flores, J. L.; Sarma, S. S. S.; López Rocha, A. N.; Nandini, S. Effects of Cladoceran-Conditioned Medium on the Demography of Brachionid Rotifers (Rotifera: Brachionidae). *Hydrobiologia* **2018**, *844* (1), 21–30. <https://doi.org/10.1007/s10750-018-3853-z>.
- Uye, S. ichi. Replacement of Large Copepods by Small Ones with Eutrophication of Embayments: Cause and Consequence. *Hydrobiologia* **1994**, *292–293* (1), 513–519.

- 7 Basterrechea-Díaz, M. *El Lago de Amatitlán: Década de Estudios Limnológicos 1985-1995*; Academia de Ciencias Médicas, Físicas y Naturales de Guatemala, Guatemala, 1997; 45 p. 308
309
- 8 Sigui, N. ¿Por Qué Continúa La Contaminación de Aguas En Guatemala? *Ciencia, Tecnol. y Salud* **2016**, *3* (2), 167–176. 310
- 9 Rodas-Pernillo, E.; Vasquez-Moscoso, C. A.; García, O. F. Dinámica Del Consumo y Aporte de Nutrientes de Fitoplancton, Dominado Por *Microcystis* sp. (Cyanophyceae) Del Lago de Amatitlán. *Ciencia, Tecnol. y Salud* **2020**, *7*, 2409–3459. 311
312
- 10 Richardson-Varas, R.; Muñoz-Luza, M.; Landeros-Cáceres, F.; Contreras-Celis, V.; Carranza-González, J.; Nuñez, M.; Hernández, E.; Alfonso-Álvarez, R.; Ramos, E.; Mazariegos, J.; Girón, J.; Velázquez, M. Índice de Fragilidad Ambiental En Las Cuencas Hidro-Geomorfológicas Del Lago Peñuelas, Chile y Del Lago Amatitlán, Guatemala. *Rev. Geográfica* **2015**, *156*, 98–109. 313
314
315
316
- 11 Brandorff, G. O. Distribution of Some Calanoida (Crustacea: Copepoda) from the Yucatán Peninsula, Belize and Guatemala. *Rev. Biol. Trop.* **2012**, *60* (1), 187–202. 317
318
- 12 Cervantes-Martínez, A.; Elías-Gutiérrez, M.; Suárez-Morales, E. Limnological and Morphometrical Data of Eight Karstic Systems “cenotes” of the Yucatan Peninsula, Mexico, during the Dry Season (February-May, 2001). *Hydrobiologia* **2002**, *482*, 167–177. 319
320
321
- 13 Schmitter-Soto, J. J.; Comín, F. A.; Escobar-Briones, E.; Herrera-Silveira, J.; Alcocer, J.; Suárez-Morales, E.; Elías-Gutiérrez, M.; Díaz-Arce, V.; Marín, L. E.; Steinich, B. Hydrogeochemical and Biological Characteristics of Cenotes in the Yucatan Peninsula (SE Mexico). *Hydrobiologia* **2002**, *467*, 215–228. 322
323
324
- 14 Jiménez, S.; Juárez, J.; Trujillo, L.; Dubón, S.; Valenzuela, O.; M, C. A. *Calidad de Agua de La Cuenca y Lago de Amatitlán*; División de Control, Calidad Ambiental y Manejo de Lagos, Guatemala, 2015; 32 p. 325
326
- 15 Elías-Gutiérrez, M.; Kotov, A. A.; Garfias-Espejo, T. Cladocera (Crustacea: Ctenopoda, Anomopoda) from Southern Mexico, Belize and Northern Guatemala, with Some Biogeographical Notes. *Zootaxa* **2006**, No. 1119, 1–27. 327
328
- 16 García-Morales, A. E.; Elías-Gutiérrez, M. The Rotifer Fauna of Guatemala and Belize: Survey and Biogeographical Affinities. *Rev. Biol. Trop.* **2007**, *55*, 569–584. 329
330
- 17 Pérez, L.; Bugja, R.; Lorenschat, J.; Brenner, M.; Curtis, J.; Hoelzmann, P.; Islebe, G.; Scharf, B.; Schwalb, A. Aquatic Ecosystems of the Yucatán Peninsula (Mexico), Belize, and Guatemala. *Hydrobiologia* **2011**, *661* (1), 407–433. 331
332
- 18 Cervantes-Martínez, A.; Elías-Gutiérrez, M.; Gutiérrez-Aguirre, M. A.; Kotov, A. A. Ecological Remarks on *Mastigodiatomus Nesus* Bowman, 1986 (Copepoda: Calanoida) in a Mexican Karstic Sinkhole. *Hydrobiologia* **2005**, *542*, 95–102. 333
334
- 19 Ramírez García, P.; Nandini, S.; Sarma, S. S. S.; Robles Valderrama, E.; Cuesta, I.; Hurtado, M. D. Seasonal Variations of Zooplankton Abundance in the Freshwater Reservoir Valle de Bravo (Mexico). *Hydrobiologia* **2002**, *467*, 99–108. 335
336
- 20 Koste, W. *Rotatoria: Die Rädertiere Mitteleuropas: Ein Bestimmunswerk Begründet von Max Voigt Überordnung Monogononta.*; Gebrüder Borntraeger, Berlin, Stuttgart, 1978; 234 p. 337
338
- 21 Fontaneto, D.; De Smet, W. H. Rotifera. In *Handbook of zoology Gastrotricha, Cycloneuralia and Gnathifera*. Vol 3; Rhaera, A. S., Ed.; De Gruyter-GmbH, Berlin, 2015; pp 217–196. 339
340
- 22 Elías-Gutiérrez, M.; Suárez-Morales, E.; Gutiérrez-Aguirre, M. A.; Silva-Briano, M.; Granados-Ramírez, J. G y Garfias-Espejo, T. *Cladocera y Copepoda de Las Aguas Continentales de México. Guía Ilustrada*; Universidad Autónoma de México, Ciudad de México, México, 2008; 322 p. 341
342
343
- 23 Suarez-Morales, E.; Gutiérrez-Aguirre, M. A.; Gómez, S.; Perbiche-Neves, G.; Previattelli, D.; Dos Santos-Silva, N.; da Rocha, C. E. F.; Mercado-Salas, N. F.; Manriquez, T. M.; Cruz-Quintana, Y.; Santana-Piñeros, A. M. Class Copepoda. In *Thorp and Covich's Freshwater Invertebrates : Volume 5: Keys to Neotropical and Antarctic Fauna*; Damborenea, C., Rogers, C. D., James, T., Eds.; Elsevier Science Publishing Co Inc., San Diego, United States, 2020; pp 663-796. 344
345
346
347
- 24 Juday, C. Limnological Studies on Some Lakes in Central America. *Wisconsin Acad. Sci. Arts, Lett.* **1915**, *18*, 214–250. 348
- 25 Wilson, M. S. New Species and Distribution Records of Diaptomid Copepods from the Marsh Collection in the United States 349

- National Museum. *J. Washingt. Acad. Sci.* **1941**, *31* (12), 509–515. 350
- 26 Brezonik, P. L.; Fox, J. L. The Limnology of Selected Guatemalan Lakes. *Hydrobiologia* **1974**, *45* (4), 467–487. 351
- 27 Ellenberg, R. L. Limnology of Lake Amatitlán in Guatemala and Its Eutrophication Process, PhD thesis, Technical University of Berlin. Berlin, Germany 128 p., 2014. 352
353
- 28 Frutos, S. M.; Poi, A. S. G.; Neiff, J. J. Zooplankton Abundance and Species Diversity in Two Lakes with Different Trophic States (Corrientes, Argentina) Abundância e Diversidade Específica Do Zooplâncton Em Dois Lagos. *Acta Limnol Bras* **2009**, *21* (3), 367–375. 354
355
356
- 29 Moreno-Gutiérrez, R. M.; Sarma, S. S. S.; Sobrino-Figueroa, A. S.; Nandini, S. Population Growth Potential of Rotifers from a High Altitude Eutrophic Waterbody, Madín Reservoir (State of Mexico, Mexico): The Importance of Seasonal Sampling. *J. Med. Virol.* **2020**, No. ii, 0–3. <https://doi.org/10.4081/jlimnol.2018>. 357
358
359
- 30 Obertegger, U.; Smith, H. A.; Flaim, G.; Wallace, R. L. Using the Guild Ratio to Characterize Pelagic Rotifer Communities. *Hydrobiologia* **2011**, *662* (1), 157–162. 360
361
- 31 Reid, J. W. *Arctodiaptomus dorsalis* (Marsh): A Case History of Copepod Dispersal. *Banisteria* **2007**, *1860* (30), 3–18. 362
- 32 Papa, R. D. S.; Li, H.; Tordesillas, D. T.; Han, B.; Dumont, H. J. Massive Invasion of *Arctodiaptomus dorsalis* (Copepoda, Calanoida, Diaptomidae) in Philippine Lakes: A Threat to Asian Zooplankton Biodiversity? *Biol. Invasions* **2012**, *14* (12), 2471–2478. 363
364
365
- 33 Metillo, E. B.; Masorong, A. M.; Macabangkit, S. A. N.; Licayan, J. R. U.; Tordesillas, D. T.; Papa, R. D. S. First Record of the Invasive *Arctodiaptomus dorsalis* (Marsh, 1907) (Copepoda: Calanoida: Diaptomidae) in Lake Lanao (Mindanao Is., Philippines). *Acta Manila. Ser. A.* **2014**, *62*, 19–23. 366
367
368
- 34 Perbiche-Neves, G.; Previattelli, D.; Pie, M. R.; Duran, A.; Suárez-Morales, E.; Boxshall, G. A.; Nogueira, M. G.; da Rocha, C. E. F. Historical Biogeography of the Neotropical Diaptomidae (Crustacea: Copepoda). *Front. Zool.* **2014**, *11* (36). <https://doi.org/10.1186/1742-9994-11-36>. 369
370
371
- 35 Elías-Gutiérrez, M.; Jerónimo, F. M.; Ivanova, N. V.; Valdez-Moreno, M.; Hebert, P. D. N. DNA Barcodes for Cladocera and Copepoda from Mexico and Guatemala, Highlights and New Discoveries. *Zootaxa* **2008**, No. 1839, 1–42. 372
373
- 36 Gutiérrez-Aguirre, M. A.; Cervantes-Martínez, A.; Elías-Gutiérrez, M.; Lugo-Vázquez, A. Remarks on *Mastigodiptomus* (Calanoida: Diaptomidae) from Mexico Using Integrative Taxonomy, with a Key of Identification and Three New Species. *PeerJ* **2020**, No. 1. <https://doi.org/10.7717/peerj.8416>. 374
375
376
- 37 Gutiérrez-Aguirre, M. A.; Suárez-Morales, E. The Eurasian *Thermocyclops crassus* (Fischer, 1853) (Copepoda, Cyclopoida) Found in Southeastern Mexico. *Crustaceana* **2000**, *73* (6), 705–713. 377
378
- 38 Gutiérrez-Aguirre, M. A.; Reid, J. W.; Suárez-Morales, E. An Afro-Asian Species of *Mesocyclops* (Copepoda: Cyclopoida) in Central America and Mexico. *J. Crustac. Biol.* **2003**, *23* (2), 352–363. 379
380
- 39 Collado, C.; Defaye, D.; Dussart, B. H.; Fernando, C. H. The Freshwater Copepoda (Crustacea) of Costa Rica with Notes on Some Species. *Hydrobiologia* **1984**, *119* (2), 89–99. 381
382
- 40 Gutiérrez-Aguirre, M. A.; Suárez-Morales, E.; Cervantes-Martínez, A.; Elías-Gutiérrez, M.; Previattelli, D. The Neotropical Species of *Mesocyclops* (Copepoda, Cyclopoida): An Upgraded Identification Key and Comments on Selected Taxa. *J. Nat. Hist.* **2006**, *40* (9–10), 549–570. 383
384
385
- 41 Suárez-Morales, E.; Gutiérrez-Aguirre, M. A.; Elías-Gutiérrez, M. Observations on the Structure of Mandibular Gnathobase in Some American *Mesocyclops* (Copepoda: Cyclopidae). *Proceedings of the Biological Society of Washington.* 2003, pp 742–753. 386
387
- 42 Cervantes-Martínez, A.; Gutiérrez-Aguirre, M. A.; Delgado-Blas, V. H.; Ruíz-Ramírez, J.-D. *Especies de Zooplancton Dulceacuicola de Cozumel*, Quintana Roo, Universidad de Quintana Roo (UQROO), Cozumel, Chetumal, 2018; 86 p. 388
389
- 43 Connolly, J. K.; Watkins, J. M.; Hinchey, E. K.; Rudstam, L. G.; Reid, J. W. New Cyclopoid Copepod (*Thermocyclops crassus*) Reported in the Laurentian Great Lakes. *J. Great Lakes Res.* **2017**, *43* (3), 198–203. 390
391

-
- 44 Verbitsky, V. B.; Lazareva, V. I.; Medyantseva, E. N.; Malysheva, O. A.; Zhdanova, S. M.; Verbitskaya, T. I.; Grishanin, A. K. 392
The Preferred and Avoidance Temperatures of *Thermocyclops crassus* (Fischer, 1853) and Their Relation to the Temperature 393
of Optimal, Pessimal and Normal Performance of the Species. *J. Therm. Biol.* **2018**, *78*, 106–113. 394